

Additive technology improves alternative feed ingredient performance

Feed additive producers are researching ways to help poultry producers deal with the challenges of using high-fiber raw materials


The use of alternative raw materials, to reduce reliance on imported materials and improve sustainability, has been much discussed. Food security is high on the agenda for government and retailers, so research into home-grown crops and co-products is important. There are economic, logistical and agronomic reasons why these resources are not used to their full potential. However, this article will look at performance limitations and how various technologies can improve them.

Challenging materials

There are several challenges to formulating with alternative raw materials. They may be higher in fiber or include anti-nutritional factors (ANF). In particular, there are higher levels of insoluble fiber in crops like sunflower and oilseed rape (canola). Arabinoxylans form complexes with protein, reducing nutrient availability. Digestibility is therefore lower and results in more undigested material in the digestive tract. This, in turn, can encourage the proliferation of pathogens, which has a negative effect on poultry health and performance.

Consistency can be a particular issue when

FIGURE 1: Improvement of ileal digestible energy from starch, fat and protein (kcal/kg DM)



Romero et al, 2014

Contribution of protein, starch and fat to the apparent ileal digestible energy of corn- and wheat-based broiler diets in response to exogenous xylanase and amylase without or with protease.

using co-products; nutrient profile and digestibility may vary between sources — and even batches. In a recent experiment, four batches of dried distillers grains with solubles (DDGS) were each fed at two different levels to laying hens. There were significant differences in digestible energy and protein digestibility between the batches that were not correlated to

differences in proximate nutrient composition. This unpredictability will have a corresponding effect on the productivity of the birds.

Processing for improvements

There are various options to process or treat raw materials prior to feeding in order to improve their nutritional value. Pulses may benefit from de-hulling to reduce the fiber and ANF levels. Heat treatment has been used to improve digestibility of a variety of cereals and pulses. Micronization, in particular, has a positive effect on insoluble fiber, meaning that nutrients have a longer time to be digested and absorbed.

The use of enzyme pre-treatment for co-products like DDGS and rapeseed meal has been investigated. The use of proteases is particularly interesting to make protein more

This is particularly true for the disruption of protein-fiber interactions, which are important in DDGS, rapeseed and sunflower.

A recent broiler digestibility study looked at the effects of adding high-fiber ingredients to corn- and wheat-based diets. Corn DDGS and rapeseed meal had the effect of reducing the availability of protein in particular. Greater effects of enzymes were found in the higher-fiber diets, increasing the combined energy contribution of protein, starch and fat (see Figure 1).

“The additive protein digestibility effect of protease on top of xylanase and amylase was also demonstrated,” Romero reiterated.

The lower digestibility of alternatives means there is the potential for more undigested material in the poultry digestive tract. This creates an environment that is favorable

for pathogen growth, making birds more susceptible to enteric disease.

Xylanase is able to create

the prebiotic compounds arabino-xylo-oligosaccharides from cereals. They encourage the growth of beneficial bacteria and the production of short-chain fatty acids.

RELATED STORY: Amino acids decrease the cost of pig feed, www.WATTAgNet.com/169013.html

digestible before it is fed. Phytases are already used in ethanol plants, resulting in lower levels of phytate in DDGS. However, cost and logistics preclude these kinds of processing despite the significant opportunities.

Enzyme technology

Enzyme technologies can help to improve feed formulation precision, diet efficiency and ultimately performance by reducing raw material variability. They also have the potential to complement endogenous enzyme production, sparing energy. When formulating with alternative raw materials, there is more potential for enzymes to have a positive effect than with standard diets.


Luis Romero, Danisco Animal Nutrition stated, “In general, you see bigger energy and protein digestibility improvements with alternatives.”

Performance benefits

Several investigations have been carried out to demonstrate the efficacy of enzymes in diets containing alternative raw materials. These recent examples were presented at the XIVth European Poultry Conference 2014 and are summarized below.

In two trials, broilers were fed diets made up of corn, corn DDGS, soybean and rapeseed meal – supplemented with either xylanase or xylanase and beta-gluconase. The enzyme combination significantly improved FCR and ileal digestibility energy, compared to the control. Both treatments significantly improved starch digestibility and tended to

Additive technology



Enzyme combinations significantly improved FCR and ileal digestibility energy in two 21-day trials, where broilers were fed diets made up of corn, corn DDGS, soybeans and rapeseed meal supplemented with either xylanase or xylanase and beta-glucanase against control.

improve ileal fat digestibility (see Figure 2).

Another study fed broilers diets containing three different levels of rapeseed and sunflower meal; with or without an enzyme combination, i.e. xylanase and beta-glucanase. Birds fed diets with high (H) levels had lower weight gain and poorer FCR, compared to those fed the medium (M) or low (L) treatments. Enzyme supplementation was shown to significantly improve FCR for the three diet types, with the highest response being seen in the H treatment (see Table 1).

Novel strategies

The combination of different groups of feed additives with potentially complementary modes of action, e.g. probiotics and enzymes, has also been investigated. These can help improve digestibility, support a healthy gut microbiota and improve bird liveability. In trials with non-challenged broilers fed a corn-soy diet containing some fibrous cereal byproducts, significant incremental increases in nitrogen corrected ap-

parent metabolizable energy (AMEn) were seen with additions of a three-strain Bacillus probiotic and xylanase, amylase and protease enzymes. Romero commented, “This combined mode of action further improves the digestibility of alternative raw materials.”

Older birds deal better with solubilized fiber due to gut maturity. This is why the use of some

TABLE 3: Broiler performance improvement (1-42 days) with and without multi-enzyme addition

Main effects	1-42 days		
	Weight gain	Feed intake	FCR
Diet type			
Low RSM	3333 ^a	5498	1.65 ^b
Low SFM/RSM	3390 ^a	5640	1.66 ^b
High SFM/RSM	3196 ^b	5486	1.72 ^a
Enzyme			
-	3290	5578	1.70 ^a
+	3323	5504	1.66 ^b
Probabilities, P <			
Diet type	**	NS	**
Enzyme	NS	NS	*
Diet type x enzyme	NS	NS	NS

NS, not significant; *, P<0.05; **, P<0.01; ***, P<0.0001.
^{a,b}Means in a column not sharing a common superscript are significantly different (P < 0.05).
¹Each value represents the mean of eight replicates (20 birds per replicate).
 RSM = Rapeseed meal; SFM = Sunflower meal
 Credit: Amerah et al, 2014

Effect of different levels of rapeseed meal or sunflower meal with or without xylanase and beta-glucanase enzyme combination on the performance of broiler chickens fed wheat-based diets. Broilers were fed diets containing three different levels of rapeseed and sunflower meal, with or without. Enzyme supplementation was shown to significantly improve FCR for the three diet types, with the highest response being seen in the High SFM/RSM treatment.

alternatives is limited or precluded in younger poultry. By having a maturation effect on the intestine, the combination of enzymes and probiotics could allow some increase in levels. Laying hens, breeders and turkeys have longer lifecycles, and so have a larger fermentative capacity and are able to better digest fiber. Therefore, there is even greater potential for enzymes to improve performance in diets containing alternative raw materials.

Future possibilities

With advancements in enzyme technology, it is possible to significantly improve poultry performance when fed rations containing alternatives. By improving the digestion and absorption of nutrients, nutritionists are able to make greater use of ingredients like DDGS, rapeseed and sunflower. This offers flexibility when formulating different types of diet, dependent on the cost and sustainability of the raw materials.

Research has shown that enzyme combinations are particularly effective, when feeding alternatives. While carbohydrases help to improve fiber digestion, proteases are interesting to get more value from lower-quality protein sources. Enzyme producers are looking at the possibility of creating raw material specific matrixes for their products. This would enable formulators to better predict the nutrient value of alternative raw materials, particularly when there is variability. Creating confidence in consistent poultry performance when fed rations containing these products. Then potentially alternative feed ingredients can be included at higher levels, benefiting efficiency and environmental impact. ◀

Zoe Kay is an independent consultant, providing technical services to the animal health and nutrition industries. She can be reached at cuckooconsulting@gmail.com.

As published in Feed International, October/November 2014